



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2008

The effect of the intergalactic environment on the observability of Ly emitters during reionization

Iliev, I T ; Shapiro, P R ; McDonald, P ; Mellema, G ; Pen, U L

Abstract: Observations of high-redshift Ly α sources are a major tool for studying the high-redshift universe and are one of the most promising ways to constrain the later stages of reionization. The understanding and interpretation of the data is far from straightforward, however. We discuss the effect of the reionizing intergalactic medium (IGM) on the observability of Ly α sources based on large simulations of early structure formation with radiative transfer. This takes into account self-consistently the reionization history, density, velocity and ionization structures and non-linear source clustering. We find that all fields are highly anisotropic and as a consequence there are very large variations in opacity among the different lines of sight. The velocity effects, from both infall and source peculiar velocity are most important for the luminous sources, affecting the line profile and depressing the bright end of the luminosity function. The line profiles are generally asymmetric and the line centres of the luminous sources are always absorbed due to the high density of the local IGM. For both luminous and average sources the damping wing effects are of similar magnitude and remain significant until fairly late, when the IGM is ionized between 30 and 70 per cent by mass. The ionizing flux in the ionized patch surrounding a high-density peak is generally strongly dominated, particularly at late times, by the cluster of faint sources, rather than the central massive galaxy. Our results reproduce well the observed mean opacity of the IGM at $z \approx 6$. The IGM absorption does not change appreciably the correlation function of sources at high redshift. Our derived luminosity function assuming constant mass-to-light ratio provides an excellent match to the shape of the observed luminosity function at $z \approx 6.6$ with faint-end slope of $\alpha = -1.5$. The resulting mass-to-light ratio implies that the majority of sources responsible for reionization are too faint to be observed by the current surveys.

DOI: <https://doi.org/10.1111/j.1365-2966.2008.13879.x>

Other titles: Effect of the intergalactic environment on the observability of Ly-alpha emitters during reionization

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-16516>

Journal Article

Accepted Version

Originally published at:

Iliev, I T; Shapiro, P R; McDonald, P; Mellema, G; Pen, U L (2008). The effect of the intergalactic environment on the observability of Ly α emitters during reionization. *Monthly Notices of the Royal Astronomical Society*, 391(1):63-83.

DOI: <https://doi.org/10.1111/j.1365-2966.2008.13879.x>

The Effect of the Intergalactic Environment on the Observability of Ly- α Emitters During Reionization

Ilian T. Iliev^{1,2*}, Paul R. Shapiro³, Patrick McDonald², Garrelt Mellema⁴, Ue-Li Pen²

¹ Universität Zürich, Institut für Theoretische Physik, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

² Canadian Institute for Theoretical Astrophysics, University of Toronto, 60 St. George Street, Toronto, ON M5S 3H8, Canada

³ Department of Astronomy, University of Texas, Austin, TX 78712-1083, U.S.A.

⁴ Stockholm Observatory, AlbaNova University Center, Stockholm University, SE-106 91 Stockholm, Sweden

DRAFT 21 October 2010 DRAFT

ABSTRACT

Observations of high-redshift Ly- α sources are a major tool for studying the high-redshift Universe and is one of the most promising ways to constrain the later stages of reionization. The understanding and interpretation of the data is far from straightforward, however. We discuss the effect of the reionizing intergalactic medium on the observability of Ly- α sources based on large simulations of early structure formation with radiative transfer. This takes into account self-consistently the reionization history, density, velocity and ionization structures and nonlinear source clustering. We find that all fields are highly anisotropic and as a consequence there are very large variations in opacity among the different lines-of-sight. The velocity effects, from both infall and source peculiar velocity are most important for the luminous sources, affecting the line profile and depressing the bright end of the luminosity function. The line profiles are generally asymmetric and the line centers of the luminous sources are always absorbed due to the high density of the local IGM. For both luminous and average sources the damping wing effects are of similar magnitude and remain significant until fairly late, when the IGM is ionized between 30% and 70% by mass.

The ionizing flux in the ionized patch surrounding a high density peak is generally strongly dominated, particularly at late times, by the cluster of faint sources, rather than the central massive galaxy. Our results reproduce well the observed mean opacity of the IGM at $z \sim 6$. The IGM absorption does not change appreciably the correlation function of sources at high redshift. Our derived luminosity function assuming constant mass-to-light ratio provides an excellent match to the shape of the observed luminosity function at $z = 6.6$ with faint-end slope of $\alpha = -1.5$. The resulting mass-to-light ratio implies that the majority of sources responsible for reionization are too faint to be observed by the current surveys.

Key words: cosmology: theory — diffuse radiation — intergalactic medium — large-scale structure of universe — galaxies: formation — radio lines: galaxies

1 INTRODUCTION

The reionization of the universe was the last global transition of the Intergalactic Medium (IGM), from fully-neutral after cosmic recombination at $z \sim 1100$ to fully-ionized as we see it today, caused by the radiation from the first stars. Currently there are still only very few direct observational constraints on this epoch. The lack of Gunn-Peterson trough in the spectra of high-redshift sources indicating a low mean neutral fraction $x_{\text{HI}} \lesssim 10^{-4}$ out to redshift $z \sim 6$, which implies overlap was achieved sometime before that, at $z_{\text{ov}} > 6$.

On the other hand, the WMAP 3-year data (Spergel & et al. 2007) yielded a fairly high value for the integrated Thomson elec-

tron scattering optical depth to the surface of last scattering, at $\tau_{\text{es}} = 0.09 \pm 0.03$. This requires a significant ionized fraction out to high redshifts, $z > 12$, and thus implies an extended reionization. The optical depth by itself does not put very stringent constraints on the possible reionization histories, however. The reason for this is the self-regulated nature of the reionization process (Iliev et al. 2007b), whereby the Jeans-mass filtering of low-mass sources in the ionized regions results in the τ_{es} and z_{ov} , the overlap redshift, being only loosely related. The overlap redshift z_{ov} is determined by the abundances and efficiencies of the high-mass sources, whose formation is not suppressed by reionization, while τ_{es} depends on both high- and low-mass sources. Thus, varying the ionizing efficiencies of the small sources yields a wide range of τ_{es} values for the same value of z_{ov} .

dramatically in the near future due to a number of large observational projects which are currently under way. The 21-cm data from high redshifts contains potentially the richest set of information since the signal is inherently three-dimensional, on the sky and in redshift/frequency (e.g. Madau et al. 1997; Tozzi et al. 2000; Iliev et al. 2002, 2003; Zaldarriaga et al. 2004; Mellema et al. 2006a; Shapiro et al. 2006; Iliev et al. 2007a, see Furlanetto et al. (2006) for a detailed recent review). The features that could be derived include the full reionization history, geometry, statistics and individual bright features (Mellema et al. 2006a; Furlanetto et al. 2006; Iliev et al. 2007a). There are significant challenges to be overcome, however, particularly related to precise subtraction of the very strong foregrounds present at low frequencies (e.g. Furlanetto et al. 2006; Morales et al. 2006; Iliev et al. 2007a).

The patchiness of reionization creates secondary temperature anisotropies in the CMB through the kinetic Sunyaev-Zel'dovich effect (Gruzinov & Hu 1998; Hu 2000; Gnedin & Jaffe 2001; Santos et al. 2003; McQuinn et al. 2005; Iliev et al. 2007c), as well as polarization anisotropies (Hu 2000; Haiman & Holder 2003; Santos et al. 2003; Mortonson & Hu 2007; Doré et al. 2007). Unlike the 21-cm signal, the reionization signatures in the CMB are integrated over the reionization history and contain no frequency information. However, the typical scales of reionization are reflected in a characteristic peak of the kSZ anisotropy signal (Iliev et al. 2007c), and the shape of the power spectrum is dependent on the reionization parameters (source efficiencies and small-scale gas clumping). CMB anisotropy observations can therefore provide us with key information about the process of reionization and since its systematics are different it would be an important complement to the 21-cm studies. One could also combine these observations more directly, by using 21-cm observations to derive the Thomson optical depth fluctuations (Holder et al. 2007). Small-scale CMB anisotropy and polarization measurements would be quite difficult due to the weakness of these signals, but are within the abilities of modern detectors (Iliev et al. 2007c; Doré et al. 2007).

Studies of the Ly- α resonant absorption in the IGM (e.g. Fan & et al. 2002; White et al. 2003; Fan & et al. 2004) and surveys of high-redshift Ly- α emitters (e.g. Rhoads & et al. 2003; Stanway & et al. 2004; Malhotra & Rhoads 2004; Bunker et al. 2006) provide important independent approaches for studying reionization (see Fan et al. 2006, for a recent review). The optical depth for Ly- α resonant absorption in neutral IGM at high redshifts is quite large (Shklovskii 1964; Gunn & Peterson 1965), thus absorption studies are mostly sensitive to very low hydrogen neutral fractions, typically below $x_{\text{HI}} \sim 10^{-4}$, otherwise the absorption saturates. This technique is thus best suited for studying highly-ionized regions and the very end of reionization and has been very successful for setting low limits for the redshift at which reionization was completed. On the other hand, Ly- α emitter surveys do not require a very low average hydrogen neutral fraction and thus in principle can probe further into the reionization epoch (Malhotra & Rhoads 2004). Narrow-band searches for high-redshift Ly- α emitters have been very successful at finding sources at ever higher redshifts, currently up to $z \sim 7$ (e.g. Hu et al. 2002, 2005; Kodaira & et al. 2003; Tran et al. 2004; Taniguchi & et al. 2005).

The correct interpretation of these data for reionization is far from straightforward, however. The strong dependence of Ly- α absorption on the neutral fraction and gas density means that both should be modelled with certain precision. High-redshift sources

generally contain multiple ionizing sources and grow much larger than the ones created by individual sources. This minimizes the effect of the damping wings and increases the transmission, allowing the detection of more and fainter sources than would be naively expected (Wyithe & Loeb 2005). However, while this is the generic expectation, the actual effect would be dependent on the exact geometry of reionization, e.g. sources close behind a neutral region will be damped even if they are inside a very large H II region. Simplified models typically assume spherical ionized regions and either ignore source clustering or assume linear bias (e.g. Cen & Haiman 2000; Santos 2004; Furlanetto et al. 2004a; Haiman & Cen 2005; Wyithe et al. 2005; Kramer et al. 2006; Wyithe & Loeb 2007). In practice the large ionized regions form by local percolation of multiple smaller ones and as a consequence are highly nonspherical (Iliev et al. 2006a; Mellema et al. 2006b; Iliev et al. 2007b). The inhomogeneous cosmological density fields and non-equilibrium chemistry effects (particularly in recently-ionized gas) further complicate the picture and point to the need of following the cosmological structure formation and the reionization history of a given region. A proper account of all these effects can only be done through detailed cosmological radiative transfer simulations.

A number of radiative transfer methods have been developed in recent years and now they are reaching a certain level of maturity and are producing fairly reliable results (Iliev & et al. 2006). However, performing large-scale reionization simulations is still technically very challenging. The majority of studies employed relatively small computational volumes and assumed optically-thin gas (e.g. Haiman & Cen 2005; Mesinger & Haiman 2007), which is the appropriate regime only well beyond the end of reionization. During reionization and shortly thereafter there are very large fluctuations in the residual neutral fraction (e.g. Wyithe & Loeb 2006) and in the corresponding optical depth, and therefore a proper radiative transfer is required for correct modelling. Recently, Maselli et al. (2007) presented a very simplified 3D treatment with radiative transfer. They considered a single luminous QSO source in a cosmological density field. They assumed that all gas starts partially-ionized to a uniform level of 90% ionized fraction and followed the evolution of the resulting H II region to derive sample absorption spectra. However, such a simple model can at best only give a rough idea about the mean radius of the resulting QSO H II region, and cannot be used to derive more detailed observational predictions. To obtain realistic spectra one has to take into account the very inhomogeneous neutral gas fraction of the IGM at $z \sim 6 - 7$, the numerous and clustered ionizing sources as well as the full, self-consistent reionization history of the region.

Kohler et al. (2005) proposed an alternative approach, whereby they used very large simulation volumes (1 Gpc to the side) to model the environments of individual, rare, luminous high-redshift quasars, utilizing results from much smaller-scale self-consistent cosmological radiative transfer simulations to calibrate the typical evolution of the coarse cells in the large box. This is currently the only viable model for simulating such rare objects, but with it one can only derive a very coarse-grained and approximate observational properties.

In this work we propose a different simulation model, whereby we use large structure formation and radiative transfer simulations to follow the full, self-consistent reionization history in a large volume of $(100 \text{ h}^{-1} \text{ Mpc})^3$ (Iliev et al. 2006b; Mellema et al. 2006a; Iliev et al. 2007b). While this volume is too small to allow us to consider the rarest, most luminous sources like the SDSS QSO's

explicit account of their radiation, and to derive good quality absorption spectra. Recently, McQuinn et al. (2007) used similar approach to study the implications of Ly- α emitters for reionization. They concentrated on the complementary topic of high-redshift Ly- α emitter surveys and what these can tell us about reionization.

Throughout this work we assume a flat ($\Omega_k = 0$) Λ CDM cosmology ($\Omega_m, \Omega_\Lambda, \Omega_b, h, \sigma_8, n$) = (0.24, 0.76, 0.042, 0.73, 0.74, 0.95) based on WMAP 3-year results (Spergel & et al. 2007), hereafter WMAP3. Here Ω_m , Ω_Λ , and Ω_b are the total matter, vacuum, and baryonic densities in units of the critical density, σ_8 is the rms density fluctuations extrapolated to the present on the scale of $8h^{-1}$ Mpc according to the linear perturbation theory, and n is the index of the primordial power spectrum of density fluctuations. In this work we consider one specific run, labelled f250C (see Iliev et al. 2007b, for details on the simulations and our notation).

2 LUMINOUS HIGH-REDSHIFT SOURCES AND THEIR ENVIRONMENT: PROPERTIES, EVOLUTION AND REIONIZATION HISTORY

The luminous sources at high redshift, the ones typically seen in current surveys, are hosted by rare, massive halos which form at the location the highest peaks of the density field. The statistics of Gaussian fields predicts that such high density peaks are rare and highly clustered in space, more strongly so at high redshifts. As a consequence, each high-redshift, massive galaxy should be surrounded by numerous smaller ionizing sources. The self-consistent reionization history simulations of such regions require following a sufficiently large volume, in order to obtain the correct statistics and biasing of the rare peaks, while at the same time resolving all the low-mass halos which are the main drivers of the reionization process. Our current radiative transfer simulations are able to achieve this. We also note that correctly modelling the nonlinear bias of the rare peaks in semi-analytical modes is a difficult and still unsolved problem. As a consequence, the halo clustering in semi-analytical models is typically underestimated, and in some cases even ignored. This often yields incorrect results, e.g. in estimates of the suppression of low-mass sources by Jeans-mass filtering (e.g. Iliev et al. 2007b).

2.1 Mass accretion history of massive halos at high redshift

In Figure 1 we show the mass-growth history of the three most massive (at redshift $z = 6$) halos found in our computational volume, with final masses of $1.5 \times 10^{12} M_\odot$, $9 \times 10^{11} M_\odot$ and $8.2 \times 10^{11} M_\odot$, respectively. All correspond to very rare, $\approx 4.5 - 5 - \sigma$ peaks of the density field. The first progenitors of these halos resolved in our simulation ($M_{\text{halo}} > 2 \times 10^9 M_\odot$) form very early, at $z \sim 16$. Thereafter, the halo masses grow roughly exponentially with redshift, $M \propto \exp(-\alpha z)$. This behaviour is in good agreement with previous results on the growth of dark matter halos in hierarchical Λ CDM models (Wechsler et al. 2002), as well as with similar results obtained in more generic, idealized situations (Shapiro et al. 2004). The slopes α for our halos exhibit much smaller scatter ($0.52 < \alpha < 0.81$; see Figure 1 caption for the complete fitting formulae) than the ones found by Wechsler et al. (2002) ($0.4 < \alpha < 1.6$). The exponential fits are excellent during




Figure 1. Mass accretion history of the three most massive halos found in our computational volume at redshift $z = 6$. The mass growth is roughly exponential in redshift, and is well-approximated by $M(z)[M_\odot] = \exp(A - \alpha z)$, where $A = 31.2$, $\alpha = 0.52$ for most massive halo (solid, red), $A = 32.5$, $\alpha = 0.81$ for the next most massive halo (short-dashed, blue) and $A = 32.0$, $\alpha = 0.75$ for the third most massive halo (long-dashed, green). Fits are shown by the thin straight lines (with corresponding line types and colors).

which time these halos have grown enough to dominate their surroundings and no similar-mass halos remain in their vicinity. The exception here is the second most massive halo (short-dashed line). Its last major merger occurs at $z \sim 8$. At early times ($z > 12$) the mass evolution of all three halos does not follow the overall exponential trend. Instead, the most massive halo is growing faster than its long-term trend, through series of rapid major mergers, while the other two halos initially grow slower than their long-term trend.

2.2 Reionization history

In Figure 2 (top and third row) we illustrate the main stages of the reionization history of a high density peak and its intergalactic environment. The particular peak we show here is the one surrounding the largest-mass halo found in our computational volume at redshift $z = 6$. For clarity and convenience for the reader we have shifted the peak to the box center using the periodicity of our computational volume. The first resolved source in this region forms at $z \sim 16$. By redshift $z = 12.9$ (top left; mass-weighted ionized fraction is $x_m = 0.001$ at this time) the central halo has already undergone several major mergers with nearby halos and its total mass has grown to $1.5 \times 10^{10} M_\odot$. The combined effect of the central halo and the nearby members of the same source cluster is to create a substantial ionized region (of size $R \sim 1 h^{-1}$ Mpc, defined as the radius at which the integrated continuum optical depth from the source at the Lyman limit reaches unity). At this time there are only a few ionized regions in our computational volume (and




Figure 2. The reionization history of a high density peak. The images are centered on the most massive (at $z = 6$) halo in our computational volume and are of size $100 h^{-1} \text{Mpc}$ to the side. The snapshots are at: (top row) $z = 12.9$ (left; global mass-weighted ionized fraction $x_m = 0.001$), $z = 10.1$ (middle; $x_m = 0.10$), $z = 9.0$ (right; $x_m = 0.28$), and (third row) $z = 7.9$ (left; $x_m = 0.66$), $z = 7.0$ (middle; $x_m = 0.94$), and $z = 6.0$ (right; $x_m = 0.9999$). The underlying cosmological density field (dark, green) is superimposed with the ionized fraction (light, orange) and the cells containing ionizing sources (dark, blue dots). The corresponding images of the (non-equilibrium) photoionization rates are shown in the second and bottom rows.




Figure 3. Histogram of the bubble size distributions along LOS. Shown are the distances from the source (the most massive halo in our volume at $z = 6$) at which continuum optical depth (at the hydrogen ionization threshold) surpasses $\tau = 1$ (black), or $\tau = 4.6$ (red), at several illustrative redshifts, as labelled.

and the ionizing regions are strongly clustered in space. The central region still remains the largest one. By redshift $z = 9.0$ (top right; $x_m = 0.28$) many more haloes have formed, most of them in large clustered groups. The H II region surrounding the central peak remains among the largest ($R \sim 6 h^{-1} \text{Mpc}$), but several other ionized bubbles reach comparable sizes.

At redshift $z = 7.9$ (bottom left; $x_m = 0.66$) some quite sizeable regions, more than ten Mpc across, have percolated, reaching local overlap. The central bubble has grown to a size of $R \sim 8 h^{-1} \text{Mpc}$. The reionization geometry becomes quite complex, with most ionized bubbles becoming interconnected, while leaving large neutral patches in-between. By $z = 7.0$ (bottom middle; $x_m = 0.94$) the notion of isolated, quasi-spherical H II regions becomes rather meaningless, since all ionized regions have already merged into one topologically-connected region, although substantial neutral patches still remain interspersed throughout our volume. The volume remains on average quite optically-thick to ionizing continuum radiation. Finally, at $z = 6.0$ (bottom right; $x_m = 0.9999$; $R = 18.7 h^{-1} \text{Mpc}$) our volume is well beyond overlap (which we define by $x_m > 99\%$). Only by that time the volume becomes on average optically-thin to ionizing radiation.

The corresponding panels in the second and fourth row of Figure 2 show images of the (non-equilibrium) ionization rate distribution at the same redshifts. The distribution is highly inhomogeneous, following the patchiness and peaking strongly in the vicinity of large source clusters. The volume-averaged photoionization rates at redshifts $z = (12.9; 10.1; 9.0; 7.9; 7.0; 6.0)$ in units of $10^{-12} s^{-1}$ are $\Gamma_{-12} = (7.0 \times 10^{-2}; 1.3 \times 10^{-1}; 2.3 \times 10^{-1}; 7.6 \times 10^{-1}; 1.3; 4.9)$, growing strongly with time as larger fraction of the volume becomes ionized and ever more sources form. Except for the earliest times, the peak photoionization rate values on the grid (corresponding to the brightest points of the images) remain

The photoionization rate distributions and evolution are discussed further in § 3.4.

To further characterize the shape and boundary sharpness of the central H II region in Figure 3 we show a histogram of the radial distance R from the central source at which the cumulative continuum optical depth at the ionizing threshold of hydrogen along each LOS reaches unity (black) and 4.6 (red; this optical depth value corresponds to 1% transmission). A spherically-symmetric H II region would correspond to a single value for the radius for a given optical depth, regardless of the directionality, while any scatter around the peak would be a measure of the non-sphericity of the ionized region. Furthermore, a comparison between the distributions for the two optical depths measures the sharpness of the H II region boundary, as follows. A sharp transition from an ionized to a neutral gas along a given LOS results in both optical depth values being surpassed simultaneously (since the neutral gas is extremely optically-thick) and in such situation the histograms would coincide. On the other hand, if the boundary of the ionized region is not clearly defined due to local percolation, gas is highly ionized and the optical depth along the LOS increases only slowly, thus the values of 1 and 4.6 are reached at very different distances from the source.

At redshift $z = 9$ the H II region surrounding the central source is largely spherical and its boundary is well-defined, albeit with some modest spread around the peak, indicating some departures from sphericity, in agreement with what was seen in Fig. 2. At redshift $z = 8$ the H II region remains fairly spherical, slightly more so for $\tau = 1$. The $\tau = 4.6$ histogram has a long tail at large values of R , up to $\sim 25 h^{-1} \text{Mpc}$, meaning that a small percentage of the LOS reach 99% opacity only at such fairly large distances. Both distributions are somewhat wider than before and start to depart from each other, reflecting the increasing “fuzziness” of the bubble boundary as it merges with other nearby bubbles due to local percolation.

At $z = 7$ the $\tau = 1$ distribution still retains a well-defined peak, albeit one accompanied by a long high- R tail reaching out to tens of Mpc. However, the $\tau = 4.6$ histogram changes its character completely, becoming very broad, with only a low peak close to the $\tau = 1$ peak and several secondary peaks. A significant number ($\sim 14\%$) of the LOS do not reach $\tau = 4.6$ within $50 h^{-1} \text{Mpc}$ of the central source (these are collected in the last, $R = 50 h^{-1} \text{Mpc}$ bin of the histogram). Finally, at $z = 6$, well beyond the overlap epoch ($z \sim 6.6$) there is no indication of a defined ionized bubble. Both distributions are very broad, with no clear peak. At this time the optical depth is determined by the tiny remaining neutral fraction, which in its turn is dictated by the density variations of the Cosmic Web. Either value of τ is reached at a wide range of distances in different directions, from $\sim 10 h^{-1} \text{Mpc}$ to over $50 h^{-1} \text{Mpc}$. The IGM becomes largely optically-thin to ionizing radiation and thus for most LOS neither $\tau = 1$ nor $\tau = 4.6$ are reached within $50 h^{-1} \text{Mpc}$ from the source.

2.3 Halo clustering in the vicinity of a luminous source

Taking a closer look at the halo clustering in the immediate vicinity of a high density peak in Fig. 4 we show the projected spatial distribution of halos around the most massive halo at $z = 7$. There are 31 resolved halos within $1 h^{-1} \text{Mpc}$ from the most massive halo and 360 resolved halos within $5 h^{-1} \text{Mpc}$ (both including the largest halo itself). The area of each circle is proportional to the mass (and




Figure 4. Projected distributions at $z = 7$ of halos within $1 h^{-1}\text{Mpc}$ comoving (left) and within $5 h^{-1}\text{Mpc}$ comoving from the most massive object in the box. Circle areas are proportional to the mass of the corresponding halo.




Figure 5. Histograms in radial bins of the total cumulative emissivity from all halos within distance R , $N_{\text{ph,tot}}(R)$, measured from the central object, here the most massive object in the computational volume, in units of the central object's emissivity, $N_{\text{ph,central}}$. Shown are (bottom to top on the right) $z = 12.9, 9, 8$ and 6.25 . The last radial bin contains the total emissivity of the computational box at that time. The vertical lines on top (corresponding line types and colors) indicate the average size of the H II region at the corresponding redshift.

centrating preferentially along the density filaments and sheets of the local Cosmic Web.

The mass of the most massive halo is $9 \times 10^{11} M_{\odot}$ at that time, well above any other halo in its vicinity. Nonetheless, the low-mass, but numerous halos surrounding the peak contribute significantly to the total ionizing emissivity. In order to quantify this point further,

of interest here. At all redshifts only within $\sim 2 h^{-1}\text{Mpc}$ from itself does the most massive halo dominate the total photoionizing emission (i.e. it contributes more than 50% of the cumulative total emission coming from that region). The exception is the highest redshift ($z = 12.9$), in which case the surrounding cluster of sources, rather than the central source, dominate the total flux even within $1 h^{-1}\text{Mpc}$ of the peak. The reason for this is the presence nearby of another halo of almost the same mass as the central one, which shortly thereafter merges with it. The cumulative emission is dominated by the small halo contribution beyond that distance. Within $10 h^{-1}\text{Mpc}$ the central source contributes only 10-30% of the total emission. As more and more halos form and the lowest mass ones become relatively common the fractional contribution of the most luminous source to the total emissivity gradually decreases in time for all radii larger than a few comoving Mpc. At $30 h^{-1}\text{Mpc}$ the central source is dominated by the rest by 1.5-2 orders of magnitude. As fraction of the total emissivity of all sources in our computational volume (shown in the last bin of the histogram) the most luminous one contributes only $\sim 2\%$ of the total at $z = 12.9$, decreasing to $\sim 0.1\%$ at $z = 6.25$.

In Figure 5 we also indicate the current H II region mean radius (vertical lines of corresponding line types and colors). Within its own bubble the central source contributes $\sim 50\%$ of the emission at $z = 12.9$, decreasing to $\sim 10\%$ at $z = 7$ and $\sim 1\%$ at $z = 6.25$.

Recently, Wyithe & Loeb (2005) presented a semi-analytical model of the source clustering at high redshift. They found that a central galaxy at $z = 7$ and with a velocity dispersion similar to our most massive source, $\sigma_V \approx 200 \text{ km s}^{-1}$) contributes $\sim 40 - 70\%$ of the H II region radius, or $\sim 10 - 35\%$ of its volume (see their Fig. 2, right panel), an over-estimate by factor of a few compared to our results where we find that the central galaxy contribution to the total flux is $\sim 1 - 10\%$. This discrepancy is most probably due to




Figure 6. Spherically-averaged profiles of (bottom to top on the left/right): the comoving number density of hydrogen (in cm^{-3} ; short-dashed, blue), the neutral hydrogen fraction (solid, red), radial velocity, v_r (in 100 km s^{-1} ; long-dashed, green) and continuum optical depth at $h\nu = 13.6 \text{ eV}$ integrated from the source ($R = 0$) outward, all at redshift $z = 6.6$ vs. R , the comoving radial distance from the most massive galaxy.




Figure 7. Mean radial velocity at $z = 6$ of the IGM with respect to the source (red solid line, negative means towards the halo, positive - away) and its rms variation (error bars), both plotted vs. (comoving) radius from the source.

2.4 IGM Environment of luminous sources at high redshift

The Ly- α absorption is strongly influenced by the local IGM environment – density, velocity and ionization structure – around the the source. As an illustrative example, let us consider a particular redshift, $z = 6.6$, similar to the highest redshifts at which currently there are Ly- α source detections. At this time the universe in our

$x_v = 0.925$. In Figure 6 we show the spherically-averaged profiles around the central source of the gas number density, n , neutral fraction, x_{HI} , integrated continuum optical depth (τ , at $h\nu = 13.6 \text{ eV}$) and radial velocity, v_r . The radiative transfer cell which contains the central source is highly-overdense, with $\delta = n/\bar{n} - 1 = 256$, which indicates that this cell roughly coincides with the source halo itself. The radial density profile declines steeply away from the source. The overdensity is $\delta = 3.3$ at $R \sim 1 \text{ h}^{-1} \text{ Mpc}$, decreasing to $\delta = 1$ at $R \sim 2.5 \text{ h}^{-1} \text{ Mpc}$, and approaching the mean density at distances beyond $R \sim 10 \text{ h}^{-1} \text{ Mpc}$.

The radial velocity profile shows an extended ($R \sim 20 \text{ h}^{-1} \text{ Mpc}$) infall region, with the mean radial velocity peaking at $\sim 150 \text{ km s}^{-1}$ before dropping to zero inside the source halo itself. The proximity region of the central source ($R < 15 \text{ h}^{-1} \text{ Mpc}$) is highly-ionized, with neutral fraction $x_{\text{HI}} < 10^{-4}$. The rest of the volume still has an appreciable neutral fraction ($\sim 0.1 - 1\%$), however, and is thus on average still optically-thick, with the mean optical depth reaching $\tau = 63$ at $R = 50 \text{ h}^{-1} \text{ Mpc}$.

However, the spherically-averaged quantities provide only a limited information about the state of the IGM surrounding each source. All quantities are distributed highly-anisotropically, and thus affect the Ly- α emission differently along each LOS. In particular, the effect of the relative velocities of the IGM and source is relatively poorly studied at present. An optically-thick medium at rest with respect to a Ly- α source would absorb the blue wing of the line and transmit its red wing, at longer wavelengths than the line centre at $\lambda_0 = 1215 \text{ \AA}$ (e.g. Miralda-Escude 1998). A relative motion of the IGM gas and the source along the LOS would result in either more or less transmission, depending on the motion direction. E.g. gas infall towards the source along the LOS from the observer would redshift it, resulting in some absorption of the red wing of the line.

In Figure 7 we show the average IGM gas velocity relative to the source (line) vs. distance from it and the variance of that average velocity (error bars). As noted above, in the vicinity of the source the IGM on average infalls towards the halo. However there are large variations, of order hundreds of km s^{-1} around this mean. E.g. a velocity offset of 200 km s^{-1} at $z = 6$ corresponds to $\Delta\lambda \sim 6 \text{ \AA}$, of the same order as the typical observed line widths, and thus a relative motion of the IGM and source of this order could have a very significant effect on the observed line. In the next section we would quantify the effect of peculiar velocities on the Ly- α line.

3 OBSERVABILITY OF HIGH- Z LY- α SOURCES

3.1 Absorption spectra of luminous sources

Much of the information about high-redshift Ly- α sources and IGM is based on absorption spectra. We thus start our discussion of the observable signatures by presenting and discussing some sample spectra intersecting the position of the most luminous source in our computational volume. Figures 8-11 we show sample spectra along three random lines-of-sight at a few selected redshifts spanning the complete range of interest here.. On the left panels we show the distributions of Ly- α Gunn-Peterson optical depth, τ_{GP} , neutral fraction, $x_{\text{HI}} = 1 - x$ (multiplied by 10^5 for clarity), and gas density in units of the mean, $\Delta = n/\bar{n}$. On the right panels we show the corresponding Gunn-Peterson transmission spectra for flux level of unity, $\exp(-\tau_{\text{GP}})$. The horizontal lines on the left panels indicate the optical depth of $\tau = 4.6$, which corresponds to 1%




Figure 8. Sample LOS at redshifts $z = 12.9$ (top) and 9.0 (bottom) vs. λ /comoving distance from the most massive galaxy. Shown are (left panels) the optical depth (solid), neutral fraction $x_{\text{HI}} = 1 - x$ ($\times 10^5$; dotted) and density in units of the mean (dashed), and (right panels) the corresponding transmission. The vertical lines show the position of the central source (in redshift space, i.e. accounting for its peculiar velocity along the LOS). The horizontal lines on the left indicate the optical depth equivalent to 1% transmission. On the right, the shaded region is the transmission in the case where the unabsorbed spectrum is flat (the horizontal dotted line).

at the mean density at redshift $z = 6.6$. All quantities shown are in redshift/wavelength space and in the observer ($z = 0$) frame. For reference, on the top axis of each figure we show the approximate corresponding distances in real space. Finally, in Figure 12 we show the mean, averaged over all random LOS, transmission spectra at the same redshifts, for the most massive source (left) and mean over all sources (right).

The Lyman- α absorption as a function of wavelength is computed using the standard procedure (e.g., Theuns et al. (1998)). The optical depth and transmission results include the redshift-space

tortions, the whole spectrum has been shifted slightly so the source is returned to its real space position). The temperature of the gas is assumed to be 10^4 K when computing thermal broadening, consistent with the assumption adopted for the simulations.

At early times ($z = 12.9$; Figure 8, top) the H II region surrounding the most massive source is still quite small (see also Figs. 2, 3 and 5) and the Ly- α emission of the source is completely suppressed by the damping wing of the Ly- α line profile, rendering it unobservable. The ionized region grows quickly after that and by $z = 9$ reaches size of $\sim 6 h^{-1} \text{Mpc}$ (Figure 8, bottom). Regardless,




Figure 9. Same as Fig. 8, but at redshifts $z = 8.1$ (top) and 7.0 (bottom).

very weak transmission. The damping wing slightly weakens compared to the higher redshifts, but is still quite substantial, and still depresses most of the red wing of the emission line.

Some of the continuum immediately behind the source in redshift space (within a few \AA) is absorbed due to gas infall towards the density peak. Because of that additional velocity towards the source the gas in front of it is slightly redshifted and thus absorbs at wavelengths red-ward of the source position. The line center at the luminous source's position is completely dark due to the high density in the middle of the density peak, regardless of the very low neutral fraction there. These features persist throughout the evolution and are very characteristic for all luminous sources, since these are always associated with high density peaks and surrounded by infall.

At redshift $z = 8.1$ (Figure 9, top) a weak damping wing is

very deep observations. Only by redshift $z = 7$ (Figures 12) the ionized region is sufficiently large for the damping wing to effectively disappear. However, the gas in the ionized region still has a significant neutral fraction, and is sufficiently dense to absorb all photons on the blue side of the Ly- α line along most LOS. On average a weak transmission at a few percent level starts coming through in the proximity region of the source (Figure 12). As more and more sources form and the ionizing flux rises the first transmission gaps start to appear at $z < 7$, both in the mean IGM away from the peak and in the source proximity (Figure 10). At redshift $z = 6.6$, our nominal overlap time (Figures 10 and 12), the proximity region extends for $\sim 30 \text{ \AA}$ and has become fairly optically-thin, allowing up to 30-40% transmission. Most of the volume still remains optically-thick, but some substantial transmission regions appear in the IGM away from the peak. We also note that there are




Figure 10. Same as Fig. 8, but at redshifts $z = 6.6$ (top) and 6.25 (bottom).

ing for much more transmission than others. The size and properties of the proximity region also vary due to its asymmetry and the anisotropy of nearby structures. Finally, during the post-overlap epoch (Figures 10, bottom, 11 and 12) the IGM slowly becomes more optically-thin to Ly- α and gradually approaches the state of the Ly- α forest. There are no more clearly-defined H II bubbles. Only a few isolated low-density regions remain neutral. This is due to the inside-out character of the reionization process, whereby the high-density regions are preferentially ionized first, while the voids, where structure formation is delayed are ionized last.

Comparing the radially-averaged mean transmission around a luminous source (high density peak) and the average for all sources (i.e. the mean behaviour around a typical source; Figure 12) we see both some similarities and several notable differences. The

($z \sim 8 - 9$) and finally disappearing at later times ($z < 7$). Naively, one might expect that compared to an average source the luminous ones would suffer from weaker damping since such sources typically reside in the middle of large H II regions, far from their boundaries. In fact, this expectation proves only partially correct. The damping is somewhat more pronounced around an average source than around a luminous one, but the differences we observe are rather modest, reflecting the fact that weaker sources are strongly clustered around the same density peaks that host the luminous ones, thus largely share the damping (or its lack) with the central source. For the same reason the damping becomes irrelevant at about the same time in both cases, which would not have been the case if the weak sources were residing in smaller, isolated bubbles.




Figure 11. Same as Fig. 8, but at redshift $z = 6.0$.




Figure 12. Mean radially-averaged transmission around the most massive source (left) and average over all sources (right) at several representative redshifts, as labelled.

of the source. The same behaviour is not seen for a typical source. Such smaller halos tend to move more in tandem with their surrounding IGM, often towards the nearest high density peak. Some local infall should exist also for these halos, but this is at very small scales, unresolved here. However, these scales are small compared to the typical emission line width (see next section) and thus we do not expect that such local infall has significant effect on the emission line.

One final important difference between a luminous and an average source is that the latter does not typically have a proxim-

transmission (10-60% transmission) regions within 5 Mpc h^{-1} ($\sim 30 \text{ \AA}$). This behaviour is again due to the high source clustering around the density peak. The combined effect of all sources is to achieve much lower neutral gas fraction regardless of the higher gas density there. The line center coinciding with a high density peak remains optically-thick, however, unlike the line center of a typical source. Away from the proximity region the absorption is largely saturated, but there are a number of transmission gaps with up to a few per cent transmission. Future work would quantify the statistics of these features and its evolution.




Figure 13. Emission lines: (left panels) intrinsic line, assumed a Gaussian with rms width of 160 km s^{-1} (black, top), and transmitted one (red, bottom) for three sample LOS at redshift $z = 8.1$, and (right panels) the corresponding optical depth (solid, black), neutral fraction $x_{\text{HI}} = 1 - x$ ($\times 10^5$; dotted, red) and density in units of the mean (dashed, green), all in redshift space (i.e. accounting for the relative velocities). The intrinsic emission line is also shown for reference.




Figure 14. Same as Fig. 13, but at redshift $z = 6.6$.

3.2 Emission line shape and its evolution

In order to study the effect of IGM absorption on the line profile shape we model the intrinsic Ly- α line as a Gaussian with an rms width of 160 km s^{-1} and peak amplitude normalized to unity. In Figures 13, 14 and 15 we show sample results for several LOS through the most luminous source at redshifts $z = 8, 6.6$ and 6 , respectively. These examples are picked to illustrate the typical cases for the observed line shape. On the left panels we show the as-

Gunn-Peterson optical depth, τ_{GP} , neutral fraction, $x_{\text{HI}} = 1 - x$ (multiplied by 10^5 for clarity), and gas density in units of the mean, $\Delta = n/\bar{n}$ and again (for reference) the intrinsic emission line.

At redshift $z = 8.1$ (Figure 13) the emission line shape is fairly regular and does not vary strongly for different LOS. The blue wing is generally highly absorbed and no appreciable flux comes through and much of the red wing is absorbed as well. The reasons for this behaviour become clear from the right panels. The neutral




Figure 15. Same as Fig. 13, but at redshift $z = 6$.




Figure 16. Evolution of the mean emission lines for most massive source (left panels) and average over all sources (right panels). Shown are the intrinsic emission line (dotted, red; assumed a Gaussian with rms width of 160 km s^{-1} , normalized to one at the peak), the transmitted line (solid, red), and mean absorption (solid, black) for three sample LOS several redshifts, as labelled.

high density of the peak and its immediate vicinity results in optical depths of $\tau_{\text{GP}} > 10$ everywhere and $\tau > 100$ at the peak itself. The gas infall towards the peak leads to a significant absorption of the red wing. The only transmitted flux comes from the far red side of the line (slightly depressed by the weak remaining damping wing).

At overlap (1% global neutral fraction by mass; $z = 6.6$) the line shape becomes much more irregular and varies significantly between the different LOS (Figure 14). Significantly larger fraction of the flux is transmitted, both on the red and on the blue side of the line. The neutral fraction in the vicinity of the source is still fairly uniform, but much lower than at the redshifts $z > 7$.

distribution. The effect of the gas infall towards the peak is still present and some of the red wing is absorbed, but much less so than at higher redshifts since the infalling gas is more highly-ionized. The line center remains absorbed and all damping wing effects have disappeared.

After overlap (Figure 15) the neutral fraction gradually declines, decreasing the optical depth and allowing ever more flux to be transmitted. The neutral fraction is still fairly uniform and thus the optical depth mostly follows the density fluctuations. The line shapes remain rather irregular, with significant variations between

In Figure 16 we show the evolution of the mean (i.e. averaged over all LOS) observed emission line shape for the most luminous source in our volume (left) and average over all sources (right). In both cases the line starts completely damped ($z = 12.9$). The later evolution of the mean line shape differs significantly, however. By redshifts $z = 9 - 10$ a significant fraction of the red wing of the line is transmitted for the luminous source, except for the line center, while much less of the red wing is transmitted for an average source. At later times ($z \sim 7 - 8$) this situation is reversed - practically all of the red wing of the line is transmitted for an average source, but much of the flux is still absorbed for the luminous source due to the high density peak in the middle and its surrounding infall. As a word of caution we should note that some of this effect is in fact not physical but numerical, since our simulations do not resolve well the detailed structure around the smaller halos. However, as we also mentioned above, these resolution effects should be modest considering that the emission line is fairly wide and thus reasonably well-resolved and any corrections due to smaller-scale structures will not affect much of the line.

On the blue side of the line there are further important differences between the luminous and average sources. The strong clustering of sources around the density peaks result in very high fluxes and thus a more pronounced highly-ionized proximity region blueward of the line center. As a consequence significantly more of the blue wing of the luminous source line is transmitted, up to 10% on average at $z = 6$, vs. only $\sim 1 - 2\%$ for an average source.

3.3 Evolution of the mean transmissivity

Figure 17 shows the mean transmission fraction as a function of redshift, for the Lyman- α , β , and γ transitions. The latter two are, respectively, 6.2 and 17.9 times weaker than Lyman- α , so transmission can be seen in cases where Lyman-alpha would be opaque (Fan & et al. 2006). At the low redshift end of our simulation data these quantities have been observed in the highest redshift quasar spectra (Fan & et al. 2006). Fig. 17 shows the (Fan & et al. 2006) (points with error bars). At $z > 6$, the Ly- α and Ly- β transmission measurements are only upper limits. It is unclear exactly what error bar should be assigned to the Ly- γ point. Fan et al. (2006) presented two upper limits and a detection in Ly- γ , which, taken together, support a detected mean transmission level (plotted in the figure) well below our prediction, with relatively small ($\sim 40\%$) errors; however, with only these three points we can not be sure that the sample variance isn't substantially larger. It appears that the reionization model in our simulations reproduces roughly the correct tail-end of reionization, but the data favor somewhat less transmission than is present in the model, i.e., a weaker radiation background and possibly slightly later reionization.

3.4 Photoionization Rates

In Figure 18 we show the normalized probability density distributions (PDFs) of the nonequilibrium photoionization rates for all cells in our computational volume at three representative redshifts - $z = 10.1$ ($x_m = 0.10$), 7.0 ($x_m = 0.94$) and 6.0 ($x_m = 0.9999$) (early times, late times and well after overlap) in linear (top) and log (bottom) scales. For comparison we also plot the photoionization rates if the corresponding cells were in ionization equilibrium. The right peak of each PDF distribution reflects the most com-




Figure 17. Overall mean transmission fraction (not necessarily near a source) in Lyman- α (solid, black), Lyman- β (dashed, red), and Lyman- γ (dotted, green). The points with horizontal error-bars show the measurements of (Fan & et al. 2006). The (black, red), (lower, upper), points with vertical error bars show Ly-(α, β), while the green dot shows Ly- γ (see text).




Figure 18. PDF of the photoionization rate at $z = 10.1$ (blue), $z = 7.0$ (red), and $z = 6.0$ (green). We show the actual, non-equilibrium rates (solid) and the corresponding equilibrium rates (dotted, same color at each redshift). All PDF's are normalized to have an area of unity below the curve.

slight shifts. The distributions are quasi-Gaussian, but with long non-Gaussian tails at both high and (especially) low values of Γ . As could have been expected, the fraction of high- Γ cells, all of which either contain sources or are in the immediate vicinity of a source, grows strongly with time, as ever. The highest photoionization rate values we find reach $\Gamma_{-12} \sim 10^3 - 10^4$. This peak value




Figure 19. Photoionization rate - overdensity correlation at $z = 9$ (left) and $z = 6$ (right). Contours are logarithmic, from 10 cells up every 0.5 dex.




Figure 20. (bottom panels) Ly- α luminosity function of high-redshift sources without (black) and with absorption included (red) at redshifts $z = 9$ (left; global mass-weighted ionized fraction $x_m = 0.28$), $z = 7$ (middle; $x_m = 0.94$) and $z = 6.0$ (right; $x_m = 0.9999$). For reference, the green, dotted line shows the result if each source is assumed 50% absorbed, which would be the case if e.g. all of the blue wing of the emission line were absorbed, while all of the red wing were transmitted. The error bar in each bin reflects the number of sources in that bin found in our computational volume. (top panels) Bin-by-bin ratio of the observed to the intrinsic luminosity function.

In the ionized regions the equilibration time is short and photoionization equilibrium is generally a good approximation. The cells with lower values of Γ (below $\Gamma_{-12} \sim 0.01 - 0.1$) correspond to the ionization fronts or neutral regions. There are many more cells in I-fronts at $z = 10.1$ than at later times, when most of the IGM is already ionized. The equilibrium and actual rates differ widely in those regions, indicating that the assumption of ionization equilibrium would be a very poor approximation there.

The photoionization rate-density correlations at $z = 9$ and $z = 6$ are shown in Figure 19 as contour plots. At high densities there is a clear, and fairly tight, correlation between the density and the photoionization rate. This reflects the fact that many more sources form in overdense regions, resulting in higher photoionization rates. At densities just above the mean the correlation becomes much less tight and around the mean densities it becomes nonexis-

(in neutral and shielded cells). There are three broad peaks of the distribution, at $\Gamma_{-12} \sim 1$ (the H II regions), $\Gamma_{-12} \sim 10^{-3}$ (cells at and around I-fronts), and $\Gamma_{-12} \sim 0$ (neutral regions). By $z = 6.0$, which is well after overlap both the neutral and self-shielded regions and the I-fronts have mostly disappeared and $\Gamma_{-12} > 0.1$ almost everywhere, rising with time as more galaxies form. These values are fairly high compared to the photoionization rate values found from the Lyman- α forest at $z \sim 2 - 4$ (Cen & McDonald 2002; Tytler & et al. 2004; Fan & et al. 2006; Bolton & Haehnelt 2007), in agreement with the high mean transmitted flux we found in § 3.3. Both point to somewhat lower ionizing source efficiencies than the ones assumed here and to a correspondingly later end of reionization.




Figure 21. Simulated luminosity function at $z = 6.6$ (dashed) vs. best fits of Kashikawa & et al. (2006) (at $z = 6.56$) for (top to bottom) faint-end slopes of $\alpha = -2, -1.5, -1$ (solid).

3.5 Luminosity function of high- z Ly- α sources

The luminosity function is an important statistical measure of the properties of high-redshift galaxies. It depends on both the intrinsic luminosity of the galaxies and the absorption in the surrounding IGM. In Figure 20 we show our results of the size of the effect of absorption on a luminosity function of high z objects. For this fiducial case we assume that the Ly- α luminosity is simply proportional to the mass of our halos (similar to our model for the ionizing sources). We compute the reduction in luminosity of each halo due to absorption (Figures 8-11 show examples of this suppression). We assume that the intrinsic Lyman- α emission line is a Gaussian with an rms of 160 km s^{-1} . Luminosity function of high-redshift sources without (in this fiducial case this is just the halo mass function; solid line) and with absorption included (dashed) at redshifts $z = 9, 7$ and 6 . For reference, the dotted line shows the result if each source is assumed 50% absorbed, which would be the case if e.g. all of the blue wing of the emission line were absorbed, while all of the red wing were transmitted. Top panels show the bin-by-bin ratios of the observed to the intrinsic luminosity function. Note that due to the binning at fixed luminosity (intrinsic or observed) this ratio is not the same as the average suppression per source of a given mass (i.e. the absorption shifts the curve both down and to the left).

The luminosity function exhibits clear evolution from high to low redshift. As we discussed in § 3.1, at high redshift the damping wings are strong and thus not only the blue side, but also significant part of the red side of the line is absorbed, as evidenced by the significant difference between the dashed and dotted lines. As a result of this absorption, the number of sources per luminosity bin drops by one to two orders of magnitude. During the later stages of reionization the damping wing effectively disappears. As a consequence, the change in the faint end of the luminosity function due to IGM absorption is on average well-represented by simply reducing each




Figure 22. Luminosity function at redshift $z=6$ if peculiar velocities are ignored (blue, long-dashed). For reference, we also show the data from Fig. 20, same notation.




Figure 23. Luminosity function at redshift $z=6$ for variable emission line width (blue, long-dashed; see text for details). For reference, we also show the data from Fig. 20, same notation.

is different at the bright end of the luminosity function, however, where on average significantly more than half of the intrinsic flux is absorbed at both $z = 7$ and $z = 6$. The shape of the luminosity function shows some evolution, as well, which is in part due to an evolution in the shape of the halo mass function and in part to the higher mean absorption for the more massive sources. The higher average absorption levels for the luminous sources is consequence of the infall which surrounds the high density peaks they are in. In

locities to zero. The result is shown in Figure 22. With no peculiar velocities present the intrinsic emission of all sources is absorbed on average at roughly the same level, by factor of 2, and the resulting luminosity function agrees well with the one where we simply assumed 50% absorption.

Our derived luminosity functions assume a constant mass-to-light ratio and are in arbitrary units (halo mass \times absorption by the IGM), proportional to a yet undetermined mass-to-observed light ratio. We can roughly determine the latter by comparing the number densities of observed and simulated objects. Kashikawa & et al. (2006) currently provide the best set of data at $z > 6$. They have provided fits to a Schechter function:

$$\phi(L)dL = \phi_* \left(\frac{L}{L_*} \right)^\alpha \exp \left(-\frac{L}{L_*} \right) \frac{dL}{L_*} \quad (1)$$

Since the high-redshift data still has large uncertainties, particularly in terms of the faint-end slope, the data is fit by assuming $\alpha = (-2, -1.5, -1)$, with best fit parameters at $z = 6.56$ given by $\log(L_*/h_{70}^{-2}\text{erg/s}) = (42.74, 42.60, 42.48)$ and $\log(\phi_*/\text{Mpc}^{-3}h_{70}^2) = (-3.14, -2.88, -2.74)$, respectively. We plot these fits in Figure 21 against our derived luminosity function. The latter was obtained by rescaling our arbitrary luminosity units to physical ones using a constant ratio,

$$L = L(M_\odot) \times 10^{30.9} \quad (2)$$

so as to match it to the observed luminosity function for the same number densities of objects.

The fit assuming $\alpha = -1.5$ provides by far the best match to our luminosity function. The two agree in both amplitude and shape over the whole available range. The differences at both the luminous and the faint end should both be expected, the former due to cosmic variance and the latter due to both numerical resolution and lack of reliable observational data. Matching the other two faint end slopes would require us to relax our constant mass-to-light ratio assumption.

Taking equation 2 at face value, we can now make an approximate correspondence between observed luminosities and masses of the underlying halos. The sources observed with Subaru at $z = 6.56$ Kashikawa & et al. (2006) have luminosities $\sim 10^{42} - 10^{42.7} \text{erg s}^{-1} h^{-2}$. This corresponds to the luminous end of our LF, for effective masses (halo mass \times absorption by the IGM) of order $10^{11} M_\odot$ or larger, or relatively rare halos. The observations are not yet sufficiently sensitive to detect the faint end, which contributes most of the ionizing emissivity during reionization. Hence, claims that observations show that there are not enough ionizing photons at $z \sim 6$ to reionize the universe appear premature.

How important is our assumption that all sources have same rms width of their intrinsic emission line profile? To check this, we replaced this assumption with one where the rms line width varies with the halo mass as $133 \text{ km/s} (M/10^{11} M_\odot)^{1/3}$ (Dijkstra et al. 2007). Results, again at $z = 6$ and in head-to-head comparison with our fiducial case of fixed line width, are shown in Figure 23. The faint end of the luminosity function proves insensitive to the line width, which is easy to understand. As we have shown above, on average the IGM completely absorbs the blue half of the line for the weaker sources and completely transmits the red half. However, the variable line width has some effect on the absorption of bright sources. Their lines become wider under this assumption and thus are less affected by the absorption due to the infalling gas, resulting in higher transmission by a factor of ~ 2 .

10 random LOS per source and we required at least 50 LOS in each bin for sampling the distribution properly. We also plot the bib-by-bin average transmission. For the luminosity bins which do not contain our minimum number of LOS we plot only the mean. Several interesting trends emerge. The distributions are fairly wide at all times and for all sources, reflecting the large variations in opacity from source to source and from LOS to LOS. The former is due to the different environments sources are found in, while the later reflects the anisotropies around each source. At all redshifts the mean and the median curves are very similar for all bins, and are essentially identical at the bright end.

The distribution itself changes its character as the evolution progresses. At early times the distribution is much wider for the fainter sources and the mean and median are gently rising towards the bright end, reflecting the fact that bright sources are found in the middle of larger H II regions, while fainter sources are found in a variety of environments. Thus, during the early evolution the main factor shaping the distribution is the local variation of the neutral fraction around each source. At late times ($z < 7$), however, the situation changes to the opposite, with the distribution becoming wider at the bright end and the mean and median decreasing there as well. By that time the IGM is already largely ionized and the main environmental dependence is due to the anisotropies of the density field and, even more so, of the infall around the bright peaks, as discussed above. This is clearly demonstrated in Fig. 25, where we show the distributions as they would be if there were no peculiar velocities. At early times the results are largely unchanged, while at late times the variations between the different LOS essentially disappear and all the curves become flat, showing that the distribution is shaped mainly by the effects of the peculiar velocities.

Finally, in Fig. 26 we show the effect of varying intrinsic line width on the distributions. Compared to our fiducial case of constant line width, at high redshift the distributions are hardly affected, except for slightly higher absorption of the faintest sources. However, at later times the varying line width has a more significant effects. At $z = 7$ the mean and the median values for the majority of sources decrease from $\sim 40 - 45\%$ down to $\sim 30\%$, but the brightest sources are affected much less. As a result the curves for the mean and median become largely flat rather than decreasing towards the bright end. Furthermore, the distribution becomes wider at the faint end, with many more LOS being absorbed by factor of 10 or more. A similar effect is seen at $z = 6$, except in this case the brightest sources are even less absorbed due to their wider emission lines, in agreement with what we observed in the luminosity functions.

4 CORRELATION FUNCTIONS

As we discussed above, the high-redshift haloes are highly clustered in space, and Ly- α sources should be clustered as well. The latter clustering has been recently observed at $z \sim 5.7$ (Murayama & et al. 2007). An interesting question to ask is if the absorption due to the surrounding IGM affects this clustering. If this were the case, then measuring the correlation function of high-redshift Ly- α sources can give us information about the state of the IGM at that time.

We derive the correlation functions as follows. First we calculate the total luminosity of each source with and without IGM




Figure 24. Transmission fraction as a function of luminosity. Shown are (dashed lines, top to bottom) 0.023, 0.16, 0.5, 0.84, and 0.977 percentiles (e.g., 2.3% of points have suppression less than the uppermost line). The center line is the median of all the LOS. We required at least 50 LOS in each bin for sampling the distribution properly. There are 10 random LOS per source. We also plot the mean (solid line) for all LOS in each bin.




Figure 25. Same as Fig. 24, but ignoring any peculiar velocities of the halos and the IGM.

by far away observer. For simplicity we consider LOS parallel to the axes of our computational box. We mock a flux-limited survey by imposing a cutoff on the observed luminosity. We compare the resulting correlation function to the one obtained for the same number of brightest sources based on their intrinsic luminosity (i.e. ignoring IGM absorption). We calculate the 3D correlation functions by direct summation over all pairs of halos as described in Martel (1991). The results at redshift $z = 9$ (with cutoff $L_{\min}(M) = 10^{9.5} M_{\odot}$) and $z = 6$ (with cutoff $L_{\min}(M) = 10^{10} M_{\odot}$) are shown in Figs. 27 and 28. In both cases the luminosity cutoffs were chosen so as to maximize the difference in the correlation functions with and without IGM absorption, while at the same time allowing for sufficient number of halos above the cutoff to reduce the noise of the correlation.

In Figure 27 we show the projection at $z = 9$ of the two source distributions onto the Y-Z plane (left), and the corresponding 3D correlation functions and their ratio (right). The projection shows that the two source populations differ, but cluster in the same spatial regions. This visual impression is confirmed by the correlation functions. We find that IGM absorption introduces only small variations in the correlation of sources. The difference is largest at

the departures are up to 5%. There are no appreciable differences at large scales.

At redshift $z = 6$ (Figure 28) there are many more sources, even with the low luminosity cutoff raised to $10^{10} M_{\odot}$. The two source distributions remain different, but are still clustered in a very similar way, resulting in largely identical correlation functions, which never differ by more than 0.2%.

The small effect of IGM absorption on the correlation function appears counter-intuitive. The reason for this is that the sources at high redshift, especially the most massive/luminous ones are strongly clustered and the ionization field is closely correlated with the galaxy field. The luminous sources are typically found in the inner parts of the largest ionized bubbles, where they are typically unaffected by damping from the remaining (few) neutral patches. This is also supported by the profiles in Fig. 12 which show the damping wing effect being important over the same time interval for both luminous and average sources. The damping affects average sources more strongly, since these are more often found closer to a neutral patch than the more massive sources. However, this does not change the correlation function significantly. Basically, the reionization patchiness has little effect on the source clustering




Figure 26. Same as Fig. 24, but for a variable Ly- α emission line width, as discussed in the text.




Figure 27. (left) Projection of the sources as seen by a mock flux-limited survey with $L > 10^{9.5} M_{\odot}$ (198 sources in total) at $z = 9$ (red points) and sources with the same number density if the IGM absorption were ignored (blue points) and (right) the 2-point 3D correlation functions of the distribution with IGM absorption (solid) and without (dashed) and their ratio (top panel).

necessarily the same individual sources) are seen as would be without IGM absorption.

At late times any clearly defined ionized regions have already disappeared and the effect of IGM absorption is to replace some sources above the luminosity cutoff with other ones essentially at random, due to small local variations of the residual neutral fraction and the gas velocities. Therefore, the reionization patchiness ultimately has little effect on the correlation function. That of course does not mean that the source population is not modified by the IGM - a flux-limited survey will see many fewer sources than if IGM absorption were not present, but the clustering properties of those sources are almost the same as with no absorption for the same number density of sources.

Recently, McQuinn et al. (2007) claimed that reionization patchiness has a significant effect on observed source clustering,

intrinsic IGM absorption for a *fixed* luminosity cutoff, rather than at fixed number density of sources, as we did. When the IGM absorption is accounted for, the sources remaining above the imposed flux limit are many fewer than in the case without absorption and they are also much brighter on average, hosted by more massive halos. This naturally results in higher bias of the observed sources compared to all sources with intrinsic luminosity above that cutoff, however, it is not necessarily related to reionization patchiness. E.g. source bias will increase also if the small residual neutral fraction in the ionized IGM increases, boosting the IGM opacity and causing dimmer, less clustered sources to fall below the luminosity cutoff, thus no neutral patches are required for this to happen. In fact, our simulation results show almost no neutral patches below $z \sim 6.6$ (at which time the global mean ionized fraction by mass is below 1%). Therefore, while our conclusions disagree with the ones of McQuinn et al. (2007), at least some of the differences could be at-




Figure 28. Projection of the sources as seen by a mock flux-limited survey with $L > 10^{10} M_{\odot}$ ($\sim 10^{40.9} h^{-2} \text{erg/s}$, according to equ. 2; 6644 sources in total) at $z = 6$ (red points) and sources with the same number density if the IGM absorption were ignored (blue points) and (right) the 2-point 3D correlation functions of the distribution with IGM absorption (solid) and without (dashed) and their ratio (top panel).

ations are real. In particular, in our simulations the pre-absorption clustering between ionized regions and sources appears to be more important than in the simulations of McQuinn et al. (2007). More detailed and direct comparisons will be required in order to evaluate and understand any differences between our results.

5 SUMMARY AND CONCLUSIONS

We considered the effects which the reionizing IGM has on the observations of high-redshift Ly- α sources. For this we utilized detailed structure formation and radiative transfer simulations, which allowed us to evaluate many features which can only be studied by detailed simulations, as well as to quantify better a number of previously-proposed effects. We followed the full reionization history self-consistently and accounted for the actual source distribution, neutral fraction, density and velocity fields.

We find that the density, neutral fraction and velocity fields are all highly anisotropic, which results in large variations in the IGM transmission and source visibility among different LOS. The velocity effects, both gas infall and source peculiar velocity are most important for massive, luminous sources. The most luminous sources are found in highest peaks of the density field, which at late times are significantly overdense out to ~ 10 comoving Mpc (cMpc) and are surrounded by infall extending to ~ 20 cMpc. The infall of gas blueshifts it in frequency space and results in significant absorption on the red side of the line center, while the peculiar velocity of the source itself can either alleviate or exacerbate this effect, depending on the halo- and infall velocity alignment.

The spherically-averaged local density enhancement and gas infall have been modelled analytically in approximate ways (Barkana 2004), and thus can be incorporated in semi-analytical models (e.g. Dijkstra et al. 2007). However, such models are unable to account for the strong intrinsic anisotropies of the neutral

tion inside the H II regions, both of which assumptions are quite unrealistic.

The Ly- α lines we derive are generally asymmetric and vary hugely from LOS to LOS. The luminous sources form at the highest density peaks and as a consequence their line centers are always highly-absorbed even though their proximity regions are very highly ionized, with typical neutral fractions $x_{\text{HI}} \sim 10^{-5} - 10^{-6}$. The luminous sources also more affected by infall and exhibit more pronounced proximity region with higher transmission of the blue wing of the line.

High-redshift sources are strongly clustered around the high peaks of the density field. The central source contributes the majority of the ionizing flux only in its immediate vicinity, within 1-2 comoving Mpc. Beyond that distance the ionizing flux is dominated by the fainter sources clustered around it. This dominance is particularly strong at late times, when both many more sources form and the ionized regions become larger, resulting in the fainter sources contributing up to 2 orders of magnitude more photons than the central source.

Compared to single-source ionized bubbles, the larger H II regions from clustered sources diminish the effects from the damping wing of the line. Nevertheless, these remain significant until fairly late (ionized mass fraction $x_m = 0.3 - 0.7$, which for the simulation considered here corresponds to redshifts $z \sim 9 - 8$). Interestingly, the average damping wing effect is similar for luminous and typical sources, even though naively one might expect that damping could be weaker for the former, since they are typically in the middle of large bubbles, away from the neutral patches, unlike the fainter sources, which are more evenly distributed.

Both the mean IGM transmission and the typical photoionization rates we find are high compared to observations at $z \sim 6$, indicating that our adopted source efficiencies are also high. The mean IGM transmissivity decreases only slowly towards the higher red-

as 1% neutral fraction) occurs at $z = 6.6$, while average neutral fraction of 10^{-4} is reached only by $z = 6$ and even then relatively small fraction (few to 10%) of the flux is transmitted. By overlap the spectra start showing significant transmission gaps in the mean IGM (i.e. away from the proximity region of a luminous source).

We find that for a given number density of sources (e.g. as determined by observations) the clustering of these sources depends only weakly on the IGM absorption during reionization. As a consequence, the reionization patchiness has little effect on the observed Ly- α source clustering, which implies that source clustering is not a good indicator for reionization patchiness.

Our derived luminosity function assuming constant mass-to-light ratio provides an excellent match to the shape of the observed luminosity function at $z = 6.6$ with faint-end slope of $\alpha = -1.5$. The resulting mass-to-light ratio implies that the majority of sources responsible for reionization are too faint to be observed by the current surveys.

ACKNOWLEDGMENTS

We thank Hugo Martel for letting us use and modify his correlation function code and X. Fan for useful discussions. This work was partially supported by NASA Astrophysical Theory Program grants NAG5-10825 and NNG04G177G to PRS.

REFERENCES

- Barkana R., 2004, MNRAS, 347, 59
 Barkana R., Loeb A., 2004, ApJ, 609, 474
 Bolton J. S., Haehnelt M. G., 2007, ArXiv Astrophysics e-prints (astro-ph/0703306)
 Bunker A., Stanway E., Ellis R., McMahon R., Eyles L., Lacy M., 2006, New Astronomy Review, 50, 94
 Cen R., Haiman Z., 2000, ApJL, 542, L75
 Cen R., McDonald P., 2002, ApJ, 570, 457
 Dijkstra M., Lidz A., Wyithe J. S. B., 2007, MNRAS, 377, 1175
 Doré O., Holder G., Alvarez M., Iliev I. T., Mellema G., Pen U.-L., Shapiro P. R., 2007, Phys. Rev. D, 76, 043002
 Fan X., Carilli C. L., Keating B., 2006, Ann. Rev. Astron. & Astrophys., 44, 415
 Fan X., et al., 2002, AJ, 123, 1247
 —, 2004, AJ, 128, 515
 —, 2006, AJ, 132, 117
 Furlanetto S. R., Hernquist L., Zaldarriaga M., 2004a, MNRAS, 354, 695
 Furlanetto S. R., Oh S. P., Briggs F. H., 2006, Physics Reports, 433, 181
 Furlanetto S. R., Zaldarriaga M., Hernquist L., 2004b, ApJ, 613, 1
 Gnedin N. Y., Jaffe A. H., 2001, ApJ, 551, 3
 Gruzinov A., Hu W., 1998, ApJ, 508, 435
 Gunn J. E., Peterson B. A., 1965, ApJ, 142, 1633
 Haiman Z., Cen R., 2005, ApJ, 623, 627
 Haiman Z., Holder G. P., 2003, ApJ, 595, 1
 Holder G. P., Iliev I. T., Mellema G., 2007, ApJL, 663, L1
 Hu E. M., Cowie L. L., Capak P., Kakazu Y., 2005, in IAU Colloq. 199: Probing Galaxies through Quasar Absorption Lines, Williams P., Shu C.-G., Menard B., eds., pp. 363–368
 Hu W., 2000, ApJ, 529, 12
 Iliev I. T., et al., 2006, MNRAS, 371, 1057
 Iliev I. T., Mellema G., Pen U.-L., Bond J. R., Shapiro P. R., 2007a, MNRAS in press, (astro-ph/0702099)
 Iliev I. T., Mellema G., Pen U.-L., Merz H., Shapiro P. R., Alvarez M. A., 2006a, MNRAS, 615
 —, 2006b, MNRAS, 369, 1625
 Iliev I. T., Mellema G., Shapiro P. R., Pen U.-L., 2007b, MNRAS, 376, 534
 Iliev I. T., Pen U.-L., Bond J. R., Mellema G., Shapiro P. R., 2007c, ApJ, 660, 933
 Iliev I. T., Scannapieco E., Martel H., Shapiro P. R., 2003, MNRAS, 341, 81
 Iliev I. T., Shapiro P. R., Ferrara A., Martel H., 2002, ApJL, 572, L123
 Kashikawa N., et al., 2006, ApJ, 637, 631
 Kodaira K., et al., 2003, Publ. Astron. Soc. Jpn., 55, L17
 Kohler K., Gnedin N. Y., Miralda-Escudé J., Shaver P. A., 2005, ApJ, 633, 552
 Kramer R. H., Haiman Z., Oh S. P., 2006, ApJ, 649, 570
 Madau P., Meiksin A., Rees M. J., 1997, ApJ, 475, 429
 Malhotra S., Rhoads J. E., 2004, ApJL, 617, L5
 Martel H., 1991, ApJ, 366, 353
 Maselli A., Gallerani S., Ferrara A., Choudhury T. R., 2007, MNRAS, 376, L34
 McQuinn M., Furlanetto S. R., Hernquist L., Zahn O., Zaldarriaga M., 2005, ApJ, 630, 643
 McQuinn M., Hernquist L., Zaldarriaga M., Dutta S., 2007, MNRAS, 381, 75
 Mellema G., Iliev I. T., Pen U.-L., Shapiro P. R., 2006a, MNRAS, 372, 679
 —, 2006b, MNRAS, 372, 679
 Mesinger A., Haiman Z., 2007, ApJ, 660, 923
 Miralda-Escudé J., 1998, ApJ, 501, 15
 Morales M. F., Bowman J. D., Hewitt J. N., 2006, ApJ, 648, 767
 Mortonson M. J., Hu W., 2007, ApJ, 657, 1
 Murayama T., et al., 2007, ApJS, 172, 523
 Rhoads J. E., et al., 2003, AJ, 125, 1006
 Santos M. G., Cooray A., Haiman Z., Knox L., Ma C.-P., 2003, ApJ, 598, 756
 Santos M. R., 2004, MNRAS, 349, 1137
 Shapiro P. R., Ahn K., Alvarez M. A., Iliev I. T., Martel H., Ryu D., 2006, ApJ, 646, 681
 Shapiro P. R., Iliev I. T., Martel H., Ahn K., Alvarez M. A., 2004, ArXiv Astrophysics e-prints (astro-ph/0409173)
 Shklovskii I. S., 1964, Astron. Zh., 41, 801
 Spergel D. N., et al., 2007, ApJS, 170, 377
 Stanway E. R., et al., 2004, ApJL, 604, L13
 Taniguchi Y., et al., 2005, Publ. Astron. Soc. Jpn., 57, 165
 Theuns T., Leonard A., Efstathiou G., Pearce F. R., Thomas P. A., 1998, MNRAS, 301, 478
 Tozzi P., Madau P., Meiksin A., Rees M. J., 2000, ApJ, 528, 597
 Tran K.-V. H., Lilly S. J., Crampton D., Brodwin M., 2004, ApJL, 612, L89
 Tytler D., et al., 2004, ApJ, 617, 1
 Wechsler R. H., Bullock J. S., Primack J. R., Kravtsov A. V., Dekel A., 2002, ApJ, 568, 52
 White R. L., Becker R. H., Fan X., Strauss M. A., 2003, AJ, 126, 1
 Wyithe J. S. B., Loeb A., 2005, ApJ, 625, 1

Wyithe J. S. B., Loeb A., Carilli C., 2005, ApJ, 628, 575

Zaldarriaga M., Furlanetto S. R., Hernquist L., 2004, ApJ, 608,
622